


## Annexe 3 de l'avis 29-2009 : fiche DDT et ses métabolites

DDT AND METABOLITES	
CAS no. 50-29-3	
Dichloro-diphenyl-trichloroethane (DDT) is an organochlorinated pesticide	
Chemical form	
	
Contamination source, origin, use	
<p>DDT was commercially introduced as an insecticide in the 1940s. Technical DDT contains 65–80 % <i>p,p'</i>-DDT. Other important constituents in the technical grade products are <i>o,p'</i>-DDT, <i>p,p'</i>-DDE and <i>p,p'</i>-DDD. The latter two compounds (along with their ortho, para analogues formed from <i>o,p'</i>-DDT) are also the major breakdown products in biological systems (EFSA, 2006). DDT and “related compounds” or sum of DDT refer to <i>p,p'</i>-DDT, <i>o,p'</i>-DDT, <i>p,p'</i>-DDE, <i>o,p'</i>-DDE, <i>p,p'</i>-DDD and <i>o,p'</i>-DDD. The main insecticidal activity can be attributed to <i>p,p'</i>-DDT (EFSA, 2006).</p> <p>DDT is used as an intermediate in the production of the pesticide dicofol and may occur as a major impurity in the final product. The use of DDT as a pesticide has been very restrictive since 1981 and banned since 1986 in the EU. Although being banned in most countries worldwide, DDT is still used for vector control especially in areas with endemic malaria, and extended use was recently recommended by WHO for indoor residual spraying to control malaria (EFSA, 2006).</p> <p>Because of the lipophilic properties and persistence in the environment, DDT and related compounds are bioaccumulated and biomagnified along the food chain. DDT is included in the Stockholm convention<sup>1</sup> on persistent organic pollutants (POPs) and the United Nations Economic Commission for Europe (UNECE) Convention on long-range transboundary air pollution protocol on POPs (CLRTAP-POP) (EFSA, 2006).</p>	
Mode of action (toxicological data: Major biologic effect, active dose)	

<sup>1</sup> The objective of this Convention is to protect human health and the environment from persistent organic pollutants.  
[http://www.pops.int/documents/convtext/convtext\\_en.pdf](http://www.pops.int/documents/convtext/convtext_en.pdf)

## **Carcinogenicity**

DDT is classified by IARC (1991) as possibly carcinogenic to humans (group 2B).

### **- Epidemiological studies**

Epidemiological studies in Colombia or in Mexico city have found a moderately high risk of breast cancer in women with higher levels of DDE (cited in Ibarluzea *et al.*, 2004).

## **Endocrine disrupting activity**

### **- Epidemiological studies**

In a study from Mexico 116 men aged 27 living in a malaria endemic area where DDT was sprayed until year 2000 in a cross-sectional study showed negative effects both on sperm motility parameters and sperm morphology which were positively correlated to plasma levels of *p,p'*-DDE (De Jager *et al.*, 2006).

### **- *in vivo* studies**

DDT increase uterine weight in rats and mice (Cited in Li and Li, 1998).

*o,p'* isomer of DDT is oestrogenic at a dose of 1 mg/kg (LOEL in rat), DDE major metabolite of DDT, has androgenic potency of about one thousandth that of dihydrotestosterone (Miyamoto and Klein, 1998).

3-Methylsulfonyl-DDE, which is a persistent but minor DDT metabolite in rats, mice and humans, is a potent adrenal toxicant in mice. This compound can be transported to the foetus through the placenta and to the offspring via mothers milk. Treatment of mice with a single dose of 3 mg/kg of 3-methylsulfonyl-DDE resulted in covalent binding of the compound to proteins followed by mitochondrial destruction in the adrenal zona fasciculata (Lund *et al.*, 1988). The binding and damage probably results from CYP11B activation in adrenal mitochondria (Jonsson *et al.*, 1991; 1995) (cited in EFSA, 2006).

### **- *in vitro* studies**

The results of competitive binding assays showed that *o,p'*-DDT, *o,p'*-DDD, *o,p'*-DDE, and *p,p'*-DDT all bind to the human estrogen receptor and *o,p'*-DDT with the strongest affinity. Binding affinities of these compounds were approximately 1000-fold weaker than that of estradiol (EFSA, 2006).

Scippo *et al.* (2004) reported even lower relative binding affinities (compared to 17 $\beta$ -estradiol) for the human estrogen receptor alpha, respectively of 0.003% (*o,p'*-DDT), 0.002% (*o,p'*-DDE and *p,p'*-DDD), 0.001% (*p,p'*-DDE), 0.0003% (*p,p'*-DDT) and 0.0006% (*o,p'*-DDD).

Higher binding affinities were reported for the human progesterone receptor, ranging from 2.5% (*p,p'*-DDT) to 0.1% (*p,p'*-DDD) (Scippo *et al.*, 2004).

#### **1. MCF-7 proliferation assays**

*o,p'*-DDT, *o,p'*-DDD, and *o,p'*-DDE were 8, 15 and 24 million times less potent than 17 $\beta$ -estradiol, in MCF-7 cells, while *o,p'*-DDT displayed one millionth the activity of 17 $\beta$ -estradiol (Cited in Dodge, 1998).

They stimulate MCF-7 cell growth at 1 $\mu$ M and 10  $\mu$ M (Cited in Li and Li, 1998).

#### **2. Transcriptional activation assays**

In transcriptional activation assays (also called reporter gene assays), *o,p'*-DDD, *p,p'*-DDD and *o,p'*-DDT have been shown to be medium agonists of the estrogen receptor, while *o,p'*-DDE and *p,p'*-DDE were slight agonist and *p,p'*-DDT displayed no agonistic activity (Willemsen *et al.*, 2004).

All DDT metabolites were potent progesterone receptor antagonists, with IC<sub>50</sub> concentrations (concentrations needed to inhibit half of the maximum response induced with progesterone) ranging between 1 to 10  $\mu$ M (Willemsen *et al.*, 2004).

The *p,p'*-DDE, hydroxy-DDE and partly *o,p'*-DDT act as anti-androgens and inhibit 5-

<p>dihydrotestosterone-induced transcriptional activation (FAO/WHO, 2001; Hartig <i>et al.</i>, 2002; Schrader and Cook, 2000) (cited in EFSA, 2006).</p> <p><i>o,p'</i>-DDT binds and activates estrogen receptors stimulating expression of other receptors for progesterone and specific enzymes and the synthesis of uterine DNA (Cited in Preziosi, 1998).</p>
<p><b>Exposure source</b></p> <p>DDT and related compounds are transferred to milk and egg and accumulate in domestic animals and fish (EFSA, 2006).</p>
<p><b>Level of exposure</b></p> <p>Mean, median and 95th percentile intakes for <i>p,p'</i>-DDT and <i>p,p'</i>-DDE were found to be 2.8, 1.5 and 6.0 and 5.3, 3.8 and 14.7 ng/kg bw per day, respectively in a German study (cited in EFSA, 2006). Based on occurrence levels in food between 1999 and 2003 and data of the Czech national consumption data base for individuals, a median dietary intake (age 4 - 90 years, both genders) of 29,1 ng/kg bw per day was calculated for total DDT (sum of <i>p,p'</i>-DDT, <i>o,p'</i>-DDT, <i>p,p'</i>-DDE, <i>o,p'</i>-DDE, <i>p,p'</i>-DDD and <i>o,p'</i>-DDD) in the Czech Republic (cited in EFSA, 2006).</p> <p>Food of animal origin is the major source of human exposure and recent studies performed in some EU Member States indicate a mean dietary intake for adults and children of 5 - 30 ng/kg bw per day. This exposure level is more than two orders of magnitude below the PTDI of 0.01 mg/kg bw (EFSA, 2006).</p> <p>Worst case exposure in Belgium (through consumption of home produced eggs, CONTEGG study): 648 ng/kg bw/day (Windal <i>et al.</i>, 2009).</p> <p>Theoretical Maximum Daily Ingestion (TMDI) in Belgium: 1290 ng/kg bw (EMRISK &amp; Ribonnet <i>et al.</i> (2007)).</p>
<p><b>TDI</b></p> <p>DDT is classified by IARC (1991) as possibly carcinogenic to humans (group 2B). The Joint FAO/WHO Meeting on Pesticide Residues (JMPR) derived a provisional tolerable daily intake (PTDI) for DDT of 0.01 mg/kg bw.</p>
<p><b>Estrogenic potency</b></p> <ul style="list-style-type: none"> <li>- Relative binding affinity Relative binding activity of DDT and related compounds is ca 10<sup>3</sup> weaker than oestradiol.</li> <li>- Uterotrophic activity <i>o,p'</i>-DDT did not affect uterus (wet and blotted) weights at doses of 100 and 500 mg/kg/day in uterotrophic assay using Sprague-Dawley immature female rats (Kim <i>et al.</i>, 2005).</li> </ul>
<p><b>% TDI</b></p> <p>The level of exposure in Europe range between 0.005 – 0.03 µg/kg bw/day (EFSA, 2006), Theoretical Maximum Daily Ingestion (TMDI) in Belgium: 1.290 µg/kg bw.</p> <p>%TDI ranges between 0.05 – 12.9%.</p>
<p><b>References</b></p> <p>De Jager C., Farias P., Barraza-Villarreal A., Hernandez Avila M., Ayotte P., Dewailly E. Dombrowski C., Rousseau, F., Diaz Sanchez V., Bailey J.L. 2006. Reduced Seminal Parameters Associated With Environmental DDT Exposure and <i>p,p'</i>-DDE Concentrations in Men in Chiapas, Mexico: A Cross-Sectional Study. <i>J. Androl.</i>, 27, 16-27.</p> <p>Dodge J.A. 1998. Natural and anthropogenic environmental oestrogens: the scientific basis for risk assessment structure/activity relationships. <i>Pure &amp; Appl. Chem.</i>, 70 (9), 1725-1733.</p>

- EFSA, 2006. Opinion of the Scientific Panel on contaminants in the food chain on a request from the commission related to DDT as an undesirable substance in animal feed. *The EFSA Journal* 433, 1 – 69.
- FAO/WHO (Food and Agriculture Organization/World Health Organization), 2001. Pesticide residues in food - 2000. Evaluations. Part II — Toxicology. Joint FAO/WHO Meeting on Pesticide Residues. <http://www.inchem.org/documents/jmpr/jmpmono/v00pr03.htm>.
- Hartig P.C., Bobseine K.L., Britt B.H., Cardon M.C., Lambright C.R., Wilson V.S., Gray L.E., Jr. 2002. Development of two androgen receptor assays using adenoviral transduction of MMTV-Luc reporter and/or hAR for endocrine screening. *Toxicol. Sci.*, 66, 82–90.
- IARC (International Agency for Research on Cancer), 1991. Occupational exposures in insecticide application, and some pesticides. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol 53, 440 pp. Lyon, France.
- Ibarluzea J. M., Fernández M. F., Santa-Marina L., Olea-Serrano M. F., Rivas A. M. Aurrekoetxea J. J. Expósito J., Lorenzo M., Torné P., Villalobos M., Pedraza V., Sasco A. J., Olea N. 2004. Breast cancer risk and the combined effect of environmental estrogens. *Cancer causes and Control*, 15, 591 – 600.
- Jonsson J., Rodriguez-Martinez H., Brandt I. 1995. Transplacental toxicity of 3-methylsulphonyl-DDE in the developing adrenal cortex in mice. *Reprod. Toxicol.*, 9(3), 257-64.
- Kim H. S., Kang T.S., Kang I.H., Kim T.S., Moon H.J., KIM I.Y., Park K.L., Lee B.M., Yoo S.D., Han S.Y. 2005. Validation study of OECD rodent uterotrophic assay for the assessment of estrogenic activity in Sprague-Dawley immature female rats. *J. Toxicol. Environ. Health A*, 68 (23-24), 2249-62.
- Li J.J., Li S.A. 1998. Natural and anthropogenic environmental oestrogens: the scientific basis for risk assessment Breast cancer: evidence for xeno-oestrogen involvement in altering its incidence and risk. *Pure & Appl. Chem.*, 70 (6), 1713-1723.
- Lund B.O., Bergman A., Brandt I. 1988. Metabolic activation and toxicity of a DDT metabolite, 3-methylsulphonyl-DDE, in the adrenal zona-fasciculata in mice. *Chem. Biol. Interact.*, 65, 25–40.
- Miyamoto J., Klein W. 1998. Natural and anthropogenic environmental oestrogens: the scientific basis for risk assessment Environmental exposure, specifies differences and risk assessment. *Pure & Appl. Chem.*, 70 (9), 1829-1845.
- Preziosi. 1998. Natural and Anthropogenic Environmental Oestrogens. The scientific basis for risk assessment. Endocrine disrupters as environmental signallers: an introduction. *Pure & Appl. Chem.*, 70 (9), 1617-1631.
- Ribonnet L., Garsou S., Pussemier L. 2007. Determination of realistic concentrations for studying toxic effects of food chemical contaminants at the gastro-intestinal level. In "Towards a Safer Food Supply in Europe", edited by C. Van Peteghem, S. De Saeger and E. Daeseleire in collaboration with L. Herman, M. De loose, E. De Pauw, P Delahaut and L. Pussemier, ISBN 978-90-8756-032-4 published in 2007 by the Belgian Science Policy, p 283-301.
- Schrader T.J., Cooke G.M. 2000. Examination of selected food additives and organochlorine food contaminants for androgenic activity in vitro. *Toxicol. Sci.*, 53 (2), 278-88.
- Scippo M.L., Argiris C., Van De Weerd C., Willemsen P., Muller M., Martial J., Maghuin-Rogister. G. 2004. Recombinant human estrogen, androgen and progesterone receptors for detection of potential endocrine disruptors. *Anal. Bioanal. Chem.*, 378, 664-669.
- Willemsen P., Scippo M.L., Kausel G., Figueroa J., Maghuin-Rogister G., Martial J.A., Muller M. 2004. Use of reporter cell lines for detection of endocrine-disrupter activity. *Anal. Bioanal. Chem.*, 378, 655-663.
- Windal I, Hanot V., Marchi J., Huysmans G., Van Overmeire I., Waegeneers N., Goeyens L. 2009. PCB and organochlorine pesticides in home-produced eggs in Belgium, *Sci. Total Environ.* (in Press), doi:10.1016/j.scitotenv.2008.11.063.